



HOW TO CARE FOR YOUR CAT FOLLOWING REPRODUCTIVE ALTERATION

PROPER RESTRAINT

Your cat may still be slightly groggy from the anesthesia as he/she leaves the office. To prevent any injury, it is best to have your cat in a pet carrier when leaving the clinic. A pet carrier will ensure the safety of your cat and decrease any chance of injury during the ride home. Upon arriving home, provide a quiet area where your cat can fully recover over the next 12-24 hours.

FOOD AND WATER

Your cat has been fasted for his/her safety in preparation for general anesthesia and surgery. Because of this, he/she may want to eat upon returning home. It is best to wait 1-2 hours after arriving home before offering a small amount of food. Offer ½ a normal meal this evening. Normal meals can be resumed tomorrow morning.

ELIMINATION

The gastrointestinal tract can be affected by general anesthesia and surgery. This may result in your cat having soft stools or constipation for 24-36 hours following the general anesthesia and surgical procedure. Some cats may have bright red blood on their stools associated with stress. If any changes persist beyond 36 hours, please contact our office.

EXERCISE AND ACTIVITY

Your cat is recovering from major surgery, so remember to limit his/her activity as much as possible. We recommend you keep him/her confined to a small area during the recovery period. **Discourage jumping of any kind.** Jumping may cause the incisions to open and require surgery to repair the damage. Normal activity may be resumed in 7-10 days.

SUTURES

Male cats do not have any sutures.

___ Absorbable sutures were used to close the incision; this will dissolve in 60-90 days.

___ Suture material was used to close the incision; the sutures need to be removed in 7-10 days.

You should check your cat's incision daily. Excessive swelling, redness or discharge should be reported immediately to our office. There should not be any bleeding. If your cat's incision begins to bleed or have any discharge, call our office immediately. Some licking is expected; if your cat begins to lick or chew obsessively at the incision sight, we can provide you with a special collar to prevent complications.

MONITOR

While we feel confident your cat will recover normally, there are risks associated with any episode of general anesthesia. Call our office if you note any abnormalities over the next week. These include, but are not limited to:

- *Loss of appetite for more than 2 days
- *Refusal to drink water for more than 1 day
- *Weakness or listlessness

*Vomiting

PAIN MEDICATION

Begin administering your cat's pain medication tomorrow morning

QUESTIONS

If you have any questions about your pet's follow-up care, please call our office immediately!